## Section V.0

Activity V.0.1 (~20 min) Consider each of the following vector properties. Label each property with  $\mathbb{R}^1$ ,  $\mathbb{R}^2$ , and/or  $\mathbb{R}^3$  if that property holds for Euclidean vectors/scalars  $\mathbf{u}, \mathbf{v}, \mathbf{w}$  of that dimension.

1. Addition associativity.

 $\mathbf{u} + (\mathbf{v} + \mathbf{w}) = (\mathbf{u} + \mathbf{v}) + \mathbf{w}.$ 

2. Addition commutivity.

 $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}.$ 

3. Addition identity.

There exists some  $\mathbf{z}$  where  $\mathbf{v} + \mathbf{z} = \mathbf{v}$ .

4. Addition inverse.

There exists some  $-\mathbf{v}$  where  $\mathbf{v} + (-\mathbf{v}) = \mathbf{z}$ .

5. Addition midpoint uniqueness.

There exists a unique  $\mathbf{m}$  where the distance from  $\mathbf{u}$  to  $\mathbf{m}$  equals the distance from  $\mathbf{m}$  to  $\mathbf{v}$ .

6. Scalar multiplication associativity.

 $a(b\mathbf{v}) = (ab)\mathbf{v}.$ 

7. Scalar multiplication identity.

 $1\mathbf{v} = \mathbf{v}.$ 

8. Scalar multiplication relativity.

There exists some scalar c where either  $c\mathbf{v} = \mathbf{w}$ or  $c\mathbf{w} = \mathbf{v}$ .

9. Scalar distribution.

 $a(\mathbf{u} + \mathbf{v}) = a\mathbf{u} + a\mathbf{v}.$ 

10. Vector distribution.

 $(a+b)\mathbf{v} = a\mathbf{v} + b\mathbf{v}.$ 

11. Orthogonality.

There exists a non-zero vector  $\mathbf{n}$  such that  $\mathbf{n}$  is orthogonal to both  $\mathbf{u}$  and  $\mathbf{v}$ .

12. Bidimensionality.

 $\mathbf{v} = a\mathbf{i} + b\mathbf{j}$  for some value of a, b.

**Definition V.0.2** A vector space V is any collection of mathematical objects with associated addition and scalar multiplication operations that satisfy the following properties. Let  $\mathbf{u}, \mathbf{v}, \mathbf{w}$  belong to V, and let a, b be scalar numbers.

- Addition associativity.
  u + (v + w) = (u + v) + w.
- Addition commutivity. u+v=v+u.
- Addition identity.

There exists some  $\mathbf{z}$  where  $\mathbf{v} + \mathbf{z} = \mathbf{v}$ .

- Addition inverse.
  - There exists some  $-\mathbf{v}$  where  $\mathbf{v} + (-\mathbf{v}) = \mathbf{z}$ .

- Scalar multiplication associativity.  $a(b\mathbf{v}) = (ab)\mathbf{v}.$
- Scalar multiplication identity.
  1v = v.
- Scalar distribution.  $a(\mathbf{u} + \mathbf{v}) = a\mathbf{u} + a\mathbf{v}.$
- Vector distribution.  $(a+b)\mathbf{v} = a\mathbf{v} + b\mathbf{v}.$

Any Euclidean vector space  $\mathbb{R}^n$  satisfies all eight requirements regardless of the value of n, but we will also study other types of vector spaces.