

Module C

Section C.1

Section C.2

Section C.3

Section C.4

Section C.5

Section C.6

Module C: Constant coefficient linear ODEs

Module C

Section C.1

Section C.2

Section C.3

Section C.4

Section C.5

Section C.6

How can we solve and apply linear constant coefficient ODEs?

Module C

Section C.1

Section C.2

Section C.3

Section C.4

Section C.5

Section C.6

At the end of this module, students will be able to...

- C1. Constant coefficient first order.** ...find the general solution to a first order constant coefficient ODE.
- C2. Modeling motion in viscous fluids.** ...model the motion of a falling object with linear drag
- C3. Homogeneous constant coefficient second order.** ...find the general solution to a homogeneous second order constant coefficient ODE.
- C4. IVPs.** ...solve initial value problems for constant coefficient ODEs
- C5. Non-homogenous constant coefficient second order.** ...find the general solution to a non-homogeneous second order constant coefficient ODE
- C6. Modeling oscillators.** ...model (free or forced, damped or undamped) mechanical oscillators with a second order ODE

Module C

Section C.1

Section C.2

Section C.3

Section C.4

Section C.5

Section C.6

Readiness Assurance Outcomes

Before beginning this module, each student should be able to...

- Describe Newton's laws in terms of differential equations.
- Find all roots of a quadratic polynomial.
- Use Euler's theorem to relate $\sin(t)$, $\cos(t)$, and e^t .
- Use Euler's theorem to simplify complex exponentials.
- Use substitution to compute indefinite integrals.
- Use integration by parts to compute indefinite integrals.
- Solve systems of two linear equations in two variables.

Module C

Section C.1

Section C.2

Section C.3

Section C.4

Section C.5

Section C.6

The following resources will help you prepare for this module.

- Describe Newtons laws in terms of differential equations.
<https://youtu.be/cioi41RrAzw>
- Find all roots of a quadratic polynomial. <https://youtu.be/2ZzuZvz33X0>
<https://youtu.be/TV5kDqiJ10s>
- Use Eulers theorem to relate $\sin(t)$, $\cos(t)$, and e^t and to simplify complex exponentials. https://youtu.be/F_0yfvm0UoU
<https://youtu.be/sn3orkHWqUQ>
- Use substitution to compute indefinite integrals.
<https://youtu.be/b76wePnIBdU>
- Use integration by parts to compute indefinite integrals.
<https://youtu.be/bZ8YAHDTFJ8>
- Solve systems of two linear equations in two variables.
<https://youtu.be/Y6JsEja15Vk>

Module C Section 1

Activity C.1.1 (~5 min)

Why don't clouds fall out of the sky?



- (a) They are lighter than air
- (b) Wind keeps them from falling
- (c) Electrostatic charge
- (d) They do fall, just very slowly

Activity C.1.2 (*~5 min*)

List all of the forces acting on a tiny droplet of water falling from the sky.

Activity C.1.3 (~ 5 min)

Tiny droplets of water obey **Stoke's law**, which says that air resistance is proportional to (the magnitude of) velocity.

- Let v be the velocity of a droplet of water (positive for upward, negative for downward).
- Let $g > 0$ be the magnitude of acceleration due to gravity and $b > 0$ be another positive constant.

Apply Newton's second law (force = mass \times acceleration) to determine which of the following **ordinary differential equations (ODEs)** models the velocity of a falling droplet of water.

(a) $v' = g - v$

(b) $v' = g + v$

(c) $mv' = -mg - bv$

(d) $mv' = -mg + bv$

Observation C.1.4

The modeling equation

$$mv' = -mg - bv$$

may be obtained by splitting the total force into gravity and air resistance:

$$F = F_g + F_r$$

Then $F = ma = mv'$ and $F_g = m(-g) = -mg$ are the result of Newton's second law, and $F_r = -bv$ holds because it should be (a) in the opposite direction of velocity and (b) a constant multiple of velocity.

Note that this equation may be rearranged as follows to group v and its derivative v' together on the left-hand side:

$$v' + \left(\frac{b}{m}\right)v = -g$$

Definition C.1.5

A **first order constant coefficient** differential equation can be written in the form

$$y' + by = f(x),$$

or equivalently,

$$\frac{dy}{dx} + by = f(x).$$


We will use both notations interchangeably.

Here, **first order** refers to the fact that the highest derivative we see is the first derivative of y .

Observation C.1.6

Consider the differential equation $y' = y$.


A useful way to visualize a first order differential equation is by a **slope field**



Each arrow represents the slope of a solution **trajectory** through that point.


Activity C.1.7 (*~5 min*)

Consider the differential equation $y' = y$ with slope field below.



Activity C.1.7 (~5 min)


Consider the differential equation $y' = y$ with slope field below.



Part 1: Draw a trajectory through the point $(0, 1)$.

Activity C.1.7 (~5 min)

Consider the differential equation $y' = y$ with slope field below.



Part 1: Draw a trajectory through the point $(0, 1)$.

Part 2: Draw a trajectory through the point $(-1, -1)$.

Activity C.1.8 (*~15 min*)

Consider the differential equation $y' = y$.

Activity C.1.8 (*~15 min*)

Consider the differential equation $y' = y$.

Part 1: Find a solution to $y' = y$.

Module C

Section C.1

Section C.2

Section C.3

Section C.4

Section C.5

Section C.6

Activity C.1.8 (*~15 min*)

Consider the differential equation $y' = y$.

Part 1: Find a solution to $y' = y$.

Part 2: Modify this solution to write an expression describing **all** solutions to $y' = y$.

Definition C.1.9

A differential equation will have many solutions. Each individual solution is said to be a **particular solution**, while the **general solution** encompasses **all** of these by using parameters such as C , k , c_0 , c_1 and so on. For example:

- The general solution to the differential equation $y' = 2x - 3$ is $y = x^2 - 3x + C$ (as done in Calculus courses).
- The general solution for $y' = y$ is $y = ke^x$ (as done in the previous activity).

Activity C.1.10 (*~15 min*)

Adapt the general solution $y = ke^x$ for $y' = y$ to find general solutions for the following differential equations.

Activity C.1.10 (*~15 min*)

Adapt the general solution $y = ke^x$ for $y' = y$ to find general solutions for the following differential equations.

Part 1: Solve $y' = 2y$.

Module C

Section C.1

Section C.2

Section C.3

Section C.4

Section C.5

Section C.6

Activity C.1.10 (*~15 min*)

Adapt the general solution $y = ke^x$ for $y' = y$ to find general solutions for the following differential equations.

Part 1: Solve $y' = 2y$.

Part 2: Solve $y' = y + 2$.

Activity C.1.11 (*~15 min*)

Find the solution for $y' = y + 2$ directly.

Simple idea: Since $y_0 = e^x$ was a particular solution of $y' = y$, we guess that a particular solution for $y' = y + 2$ is of the form $y_p = ve^x$ for some **function** $v(x)$.

Activity C.1.11 (*~15 min*)

Find the solution for $y' = y + 2$ directly.

Simple idea: Since $y_0 = e^x$ was a particular solution of $y' = y$, we guess that a particular solution for $y' = y + 2$ is of the form $y_p = ve^x$ for some **function** $v(x)$.

Part 1: Use the Product Rule to find $y'_p = \frac{d}{dx}[ve^x]$.

Activity C.1.11 (*~15 min*)

Find the solution for $y' = y + 2$ directly.

Simple idea: Since $y_0 = e^x$ was a particular solution of $y' = y$, we guess that a particular solution for $y' = y + 2$ is of the form $y_p = ve^x$ for some **function** $v(x)$.

Part 1: Use the Product Rule to find $y'_p = \frac{d}{dx}[ve^x]$.

Part 2: Substitute y_p and y'_p into the equation $y' = y + 2$.

Activity C.1.11 (*~15 min*)

Find the solution for $y' = y + 2$ directly.

Simple idea: Since $y_0 = e^x$ was a particular solution of $y' = y$, we guess that a particular solution for $y' = y + 2$ is of the form $y_p = ve^x$ for some **function** $v(x)$.

Part 1: Use the Product Rule to find $y'_p = \frac{d}{dx}[ve^x]$.

Part 2: Substitute y_p and y'_p into the equation $y' = y + 2$.

Part 3: Solve for v' , and integrate to find v .

Activity C.1.11 (*~15 min*)

Find the solution for $y' = y + 2$ directly.

Simple idea: Since $y_0 = e^x$ was a particular solution of $y' = y$, we guess that a particular solution for $y' = y + 2$ is of the form $y_p = ve^x$ for some **function** $v(x)$.

Part 1: Use the Product Rule to find $y'_p = \frac{d}{dx}[ve^x]$.

Part 2: Substitute y_p and y'_p into the equation $y' = y + 2$.

Part 3: Solve for v' , and integrate to find v .

Part 4: Find y_p .

Observation C.1.12

The technique outlined in the previous activity is called **variation of parameters**. If y_0 is a particular solution of the **homogeneous** equation, assume that a particular solution of the **non-homogeneous** equation has the form $y_p = vy_0$, and then determine what v must be.

Example:

$$y' + 3y = 0$$

homogeneous

$$y' + 3y = x$$

non-homogeneous

Note that each term of the homogeneous equation includes y or its derivatives.

Activity C.1.13 (*~20 min*)

Solve $y' = x - 3y$ by first solving its corresponding homogeneous equation and using variation of parameters:

$$y' + 3y = 0$$

homogeneous

$$y' + 3y = x$$

non-homogeneous

Activity C.1.13 (*~20 min*)

Solve $y' = x - 3y$ by first solving its corresponding homogeneous equation and using variation of parameters:

$$y' + 3y = 0$$

homogeneous

$$y' + 3y = x$$

non-homogeneous

Part 1: Modify e^x to find the general solution y_h for the homogeneous equation.

Activity C.1.13 (~ 20 min)

Solve $y' = x - 3y$ by first solving its corresponding homogeneous equation and using variation of parameters:

$$y' + 3y = 0$$

homogeneous

$$y' + 3y = x$$

non-homogeneous

Part 1: Modify e^x to find the general solution y_h for the homogeneous equation.

Part 2: Choose a particular solution y_0 for the homogeneous equation, and assume $y_p = v y_0$ is a particular solution of the non-homogeneous equation for some **function** v . Substitute y_p into non-homogeneous equation and simplify.

Activity C.1.13 (*~20 min*)

Solve $y' = x - 3y$ by first solving its corresponding homogeneous equation and using variation of parameters:

$$y' + 3y = 0$$

homogeneous

$$y' + 3y = x$$

non-homogeneous

Part 1: Modify e^x to find the general solution y_h for the homogeneous equation.

Part 2: Choose a particular solution y_0 for the homogeneous equation, and assume $y_p = v y_0$ is a particular solution of the non-homogeneous equation for some **function** v . Substitute y_p into non-homogeneous equation and simplify.

Part 3: Determine v , and then determine y_p .

Observation C.1.14

Since $y_h = ke^{-3x}$ was the general solution of $y' + 3y = 0$, and $y_p = \frac{x}{3} - \frac{1}{9}$ is a particular solution of $y' + 3y = x$,

$$y = y_h + y_p = (ke^{-3x}) + \left(\frac{x}{3} - \frac{1}{9}\right)$$

is a solution to $y' + 3y = x$:

$$\frac{d}{dx}[y_h + y_p] + 3(y_h + y_p) = (y_h' + 3y_h) + (y_p' + 3y_p) = 0 + x = x$$

Fact C.1.15

Let a be a constant real number. Every constant coefficient first order ODE

$$y' + ay = f(x)$$

has the general solution

$$y = y_h + y_p$$

where y_h is the general solution to the homogeneous equation $y' + ay = 0$ and y_p is a particular solution to $y' + ay = f(t)$.

Activity C.1.16 (*~15 min*)

Find the general solution to $y' = 2y + x + 1$ using variation of parameters:

- Write the homogeneous equation and find its general solution y_h .
- Use a particular solution y_0 for the homogeneous equation to find a particular solution $y_p = vy_0$ for the original equation.
- Then $y = y_h + y_p$ gives the general solution to the equation.

Module C Section 2

Observation C.2.1

Recall that we can model the velocity of a water droplet in a cloud by

$$mv' = -mg - bv$$

where negative numbers represent downward motion, $m > 0$ is the mass of the droplet, $g > 0$ is the magnitude of acceleration due to gravity, and $b > 0$ is the proportion of wind resistance to speed.



Activity C.2.2 (*~20 min*)

A water droplet with a radius of $10\ \mu\text{m}$ has a mass of about $4 \times 10^{-15}\ \text{kg}$. It is determined in a laboratory that for a droplet this size, the constant b has a value of $3 \times 10^{-3}\ \text{kg/s}$, and it is known that g is approximately $9.8\ \text{m/s}^2$.

Complete the following tasks to study the motion of this droplet.

Activity C.2.2 (*~20 min*)

A water droplet with a radius of $10\ \mu\text{m}$ has a mass of about 4×10^{-15} kg. It is determined in a laboratory that for a droplet this size, the constant b has a value of 3×10^{-3} kg/s, and it is known that g is approximately $9.8\ \text{m/s}^2$.

Complete the following tasks to study the motion of this droplet.

Part 1: Rewrite $mv' = -mg - bv$ in the form of $v' + av = ?$ for some value of a .

Activity C.2.2 (*~20 min*)

A water droplet with a radius of $10\ \mu\text{m}$ has a mass of about 4×10^{-15} kg. It is determined in a laboratory that for a droplet this size, the constant b has a value of 3×10^{-3} kg/s, and it is known that g is approximately $9.8\ \text{m/s}^2$.

Complete the following tasks to study the motion of this droplet.

Part 1: Rewrite $mv' = -mg - bv$ in the form of $v' + av = ?$ for some value of a .

Part 2: Find the general solution of this ODE in terms of a and g . (Let $v_p = wv_0$ when using variation of parameters to avoid confusion.)

Activity C.2.2 (~ 20 min)

A water droplet with a radius of $10\ \mu\text{m}$ has a mass of about 4×10^{-15} kg. It is determined in a laboratory that for a droplet this size, the constant b has a value of 3×10^{-3} kg/s, and it is known that g is approximately $9.8\ \text{m/s}^2$.

Complete the following tasks to study the motion of this droplet.

Part 1: Rewrite $mv' = -mg - bv$ in the form of $v' + av = ?$ for some value of a .

Part 2: Find the general solution of this ODE in terms of a and g . (Let $v_p = wv_0$ when using variation of parameters to avoid confusion.)

Part 3: Due to wind resistance, eventually the droplet will effectively stop accelerating upon reaching a certain velocity. What is this **terminal velocity** of the droplet in terms of a and g ?

Activity C.2.2 (~ 20 min)

A water droplet with a radius of $10 \mu\text{m}$ has a mass of about 4×10^{-15} kg. It is determined in a laboratory that for a droplet this size, the constant b has a value of 3×10^{-3} kg/s, and it is known that g is approximately 9.8 m/s^2 .

Complete the following tasks to study the motion of this droplet.

Part 1: Rewrite $mv' = -mg - bv$ in the form of $v' + av = ?$ for some value of a .

Part 2: Find the general solution of this ODE in terms of a and g . (Let $v_p = wv_0$ when using variation of parameters to avoid confusion.)

Part 3: Due to wind resistance, eventually the droplet will effectively stop accelerating upon reaching a certain velocity. What is this **terminal velocity** of the droplet in terms of a and g ?

Part 4: If the droplet starts from rest ($v = 0$ when $t = 0$), what is its velocity after 0.01 s? Use a calculator to compute the answer in m/s.

Definition C.2.3

The last part of the previous activity is an example of an **Initial Value Problem (IVP)**; we were given the initial value of the velocity in addition to our differential equation.

$$v' + (b/m)v = -g \quad v(0) = 0$$

Physical scenarios often produce IVPs with a unique solution.

Module C Section 3


Observation C.3.1

What happens when your tire hits a pothole?

<https://prof.clontz.org/assets/img/good-bad-shocks.gif>

Activity C.3.2 (~ 5 min)

Hooke's law says that the force exerted by the spring is proportional to the distance the spring is stretched from its natural length, given by a spring coefficient $k > 0$.




Let y measure the displacement of the mass from the spring's natural length. Write a differential equation modeling the displacement of the m kg mass, assuming that the only force acting on the mass comes from the spring.

Observation C.3.3

Since the spring acts on the mass in the opposite direction of displacement, we may model the mass-spring system with

$$my'' = -ky.$$



Module C

Section C.1

Section C.2

Section C.3

Section C.4

Section C.5

Section C.6

Activity C.3.4 (*~15 min*)

Consider the mass-spring equation $my'' = -ky$ where $m = k = 1$:

$$y'' = -y.$$

Module C

Section C.1

Section C.2

Section C.3

Section C.4

Section C.5

Section C.6

Activity C.3.4 (*~15 min*)

Consider the mass-spring equation $my'' = -ky$ where $m = k = 1$:

$$y'' = -y.$$

Part 1: Find a solution.

Module C

Section C.1

Section C.2

Section C.3

Section C.4

Section C.5

Section C.6

Activity C.3.4 (*~15 min*)

Consider the mass-spring equation $my'' = -ky$ where $m = k = 1$:

$$y'' = -y.$$

Part 1: Find a solution.

Part 2: Find the general solution.

Module C

Section C.1

Section C.2

Section C.3

Section C.4

Section C.5

Section C.6

Activity C.3.4 (*~15 min*)

Consider the mass-spring equation $my'' = -ky$ where $m = k = 1$:

$$y'' = -y.$$

Part 1: Find a solution.


Part 2: Find the general solution.

Part 3: Describe the long term behavior of the mass-spring system.

Activity C.3.5 (~ 5 min)

The general solution $y = c_1 \cos(t) + c_2 \sin(t)$ models infinitely oscillating behavior, but in applications this does not occur.

Thus, a damper (a.k.a. dashpot) is often considered, which provides a force proportional to velocity, given by the coefficient $b > 0$. For example, friction may act as a damper to a mass-spring system.



Write a differential equation modeling the displacement of a mass in a **damped** mass-spring system.

Observation C.3.6

The damped mass-spring system can be modelled by

$$my'' = -by' - ky.$$

Here m is the mass, k is the spring constant, and b is the damping constant. We can rearrange this as

$$y'' + By' + Ky = 0$$

where $B = \frac{b}{m}$ and $K = \frac{k}{m}$.

This is a **homogeneous second order constant coefficient** differential equation. Here, **homogeneous** refers to the 0 on the right hand side of the equation.

Activity C.3.7 (*~15 min*)

Consider the second order constant coefficient equation

$$y'' = y.$$

Module C

Section C.1

Section C.2

Section C.3

Section C.4

Section C.5

Section C.6

Activity C.3.7 (*~15 min*)

Consider the second order constant coefficient equation

$$y'' = y.$$

Part 1: Find a solution.

Module C

Section C.1

Section C.2

Section C.3

Section C.4

Section C.5

Section C.6

Activity C.3.7 (*~15 min*)

Consider the second order constant coefficient equation

$$y'' = y.$$

Part 1: Find a solution.*Part 2:* Find the general solution.

Module C

Section C.1

Section C.2

Section C.3

Section C.4

Section C.5

Section C.6

Activity C.3.7 (*~15 min*)

Consider the second order constant coefficient equation

$$y'' = y.$$

Part 1: Find a solution.*Part 2:* Find the general solution.*Part 3:* Describe the long term behavior of the solutions.

Observation C.3.8

It is sometimes useful to think in terms of **differential operators**.

- We will use D to represent a derivative. So for any function y ,

$$D(y) = \frac{\partial y}{\partial x} = y'.$$

- D^2 will denote the second derivative operator (i.e. differentiate twice, or apply D twice).
- We will use I for the identity operator, so $I(y) = y$. (It can be thought of as $I = D^0$, take the derivative zero times.)

In this language, the differential equation $y' + 3y = 0$ can be rewritten as $D(y) + 3I(y) = 0$, or more simply $(D + 3I)(y) = 0$.

Thus, the question of solving the homogeneous differential equation is the question of finding the **kernel** of the differential operator $D + 3I$: all the functions y that the transformation $D + 3I$ turns into the zero function.

Module C

Section C.1

Section C.2

Section C.3

Section C.4

Section C.5

Section C.6

Activity C.3.9 (*~5 min*)Find a differential operator whose kernel is the solution set of the ODE $y' = 4y$.

- a) $D - 4I$
- b) $D + 4I$
- c) $D^2 - 4I$
- d) $D^2 + 4D$

Activity C.3.10 (*~5 min*)

The kernel of the differential operator $D - 4I$ whose kernel is the general solution of the ODE $y' = 4y$. What is its general solution?

- a) $y = ke^{-4x}$
- b) $y = ke^{4x}$
- c) $y = 4x + k$
- d) $y = 4$

Activity C.3.11 (*~5 min*)

What are ODE and general solution given by the kernel of the differential operator $D - aI$ for a real number a ?

- a) $y' - ay = 0$ and $y = ke^{ax}$.
- b) $y' + ay = 0$ and $y = ke^{-ax}$.
- c) $y' - a = 0$ and $y = ax + k$.
- d) $y'' + a = 0$ and $y = -\frac{a}{2}x^2 + kx + l$.

Module C

Section C.1

Section C.2

Section C.3

Section C.4

Section C.5

Section C.6

Observation C.3.12

The kernel of the differential operator $D - aI$ is given by the general solution $y = ke^{ax}$.

Activity C.3.13 (*~15 min*)

Consider the ODE

$$y'' + 5y' + 6y = 0.$$

Activity C.3.13 (*~15 min*)

Consider the ODE

$$y'' + 5y' + 6y = 0.$$

Part 1: Use I, D, D^2 to write a differential operator whose kernel is the solution set of the above ODE.

Activity C.3.13 (*~15 min*)

Consider the ODE

$$y'' + 5y' + 6y = 0.$$

Part 1: Use I, D, D^2 to write a differential operator whose kernel is the solution set of the above ODE.

Part 2: Factor this differential operator as a composition of two simpler operators, as you would a polynomial. (This works because the order of applying the transformations D and I doesn't matter).

Activity C.3.13 (~ 15 min)

Consider the ODE

$$y'' + 5y' + 6y = 0.$$

Part 1: Use I, D, D^2 to write a differential operator whose kernel is the solution set of the above ODE.

Part 2: Factor this differential operator as a composition of two simpler operators, as you would a polynomial. (This works because the order of applying the transformations D and I doesn't matter).

Part 3: Find the general solution for each factor, and then combine to find the general solution to the overall ODE.

Activity C.3.13 (*~15 min*)

Consider the ODE

$$y'' + 5y' + 6y = 0.$$

Part 1: Use I, D, D^2 to write a differential operator whose kernel is the solution set of the above ODE.

Part 2: Factor this differential operator as a composition of two simpler operators, as you would a polynomial. (This works because the order of applying the transformations D and I doesn't matter).

Part 3: Find the general solution for each factor, and then combine to find the general solution to the overall ODE.

Part 4: Check that your general solution is valid by computing y', y'' and plugging into $y'' + 5y' + 6y = 0$.

Observation C.3.14

The kernel of $(D + 3I)(D + 2I)$ is given by $y = k_1e^{-3t} + k_2e^{-2t}$.

In general for $\alpha \neq \beta$, the kernel of $(D - \alpha I)(D - \beta I)$ is given by $y = k_1e^{at} + k_2e^{bt}$.

Module C

Section C.1

Section C.2

Section C.3

Section C.4

Section C.5

Section C.6

Activity C.3.15 (*~10 min*)

Solve the ODE

$$2y'' + 7y' + 6y = 0.$$

Module C

Section C.1

Section C.2

Section C.3

Section C.4

Section C.5

Section C.6

Activity C.3.16 (*~15 min*)

Recall that the general solution to $y'' + y = 0$ is given by $y = c_1 \sin(x) + c_2 \cos(x)$. Show how to find this solution using the differential operator $D^2 + 1$.

Module C

Section C.1

Section C.2

Section C.3

Section C.4

Section C.5

Section C.6

Activity C.3.17 (*~15 min*)

Consider the ODE

$$y'' + 2y' + 5y = 0$$

.

Module C

Section C.1

Section C.2

Section C.3

Section C.4

Section C.5

Section C.6

Activity C.3.17 (*~15 min*)

Consider the ODE

$$y'' + 2y' + 5y = 0$$

.

Part 1: Find its general solution using complex numbers.

Activity C.3.17 (*~15 min*)

Consider the ODE

$$y'' + 2y' + 5y = 0$$

.

Part 1: Find its general solution using complex numbers.*Part 2:* Describe the general solution only involving real numbers.

Activity C.3.18 (*~5 min*)

Which of these are solutions to the following ODE?

$$y'' - 4y' + 4y = 0$$

- a) $y = e^{2t}$, where $y' = 2e^{2t}$ and $y'' = 4e^{2t}$
- b) $y = te^{2t}$, where $y' = e^{2t} + 2te^{2t}$ and $y'' = 4e^{2t} + 4e^{2t}$
- c) $y = e^{2t} + te^{2t}$, where $y' = 3e^{2t} + 2te^{2t}$ and $y'' = 8e^{2t} + 4e^{2t}$
- d) All of the above

Observation C.3.19

To solve $y'' - 4y' + 4y = 0$, we need to find the kernel of $(D - 2I)(D - 2I) = (D - 2I)^2$.

- The kernel of $D - 2I$ is given by ke^{2x} .
- But if $(D - 2I)(y) = e^{2t}$, then $(D - 2I)(D - 2I)(y) = (D - 2I)(e^{2t}) = 0$ also.
- That means the kernel of $(D - 2I)^2$ is given by both $(D - 2I)(y) = 0$ and $(D - 2I)(y) = e^{2t}$.

Module C

Section C.1

Section C.2

Section C.3

Section C.4

Section C.5

Section C.6

Activity C.3.20 (*~15 min*)Solve $(D - 2I)(y) = e^{2x}$.

Observation C.3.21

Since $(D - 2I)(y) = 0$ solves to ke^{2t} and $(D - 2I)(y) = e^{2t}$ solves to kte^{2t} , we have shown that the general solution of

$$y'' - 4y' + 4y = 0$$

is

$$y = c_0e^{2t} + c_1te^{2t}.$$

Activity C.3.22 (*~10 min*)

Consider the homogeneous second order constant coefficient ODE

$$ay'' + by' + cy = 0.$$

If r is a number such that $ar^2 + br + c = 0$, what can you conclude?

- (a) e^{rt} is a solution.
- (b) e^{-rt} is a solution.
- (c) te^{rt} is a solution.
- (d) There are no solutions.

Activity C.3.23 (*~5 min*)

Consider the homogeneous second order constant coefficient ODE

$$ay'' + by' + cy = 0.$$

When does the general solution have the form $c_0e^{rt} + c_1te^{rt}$?

- (a) When the polynomial $ax^2 + bx + c$ has two distinct real roots.
- (b) When the polynomial $ax^2 + bx + c$ has a repeated real root.
- (c) When the polynomial $ax^2 + bx + c$ has two distinct non-real roots.
- (d) When the polynomial $ax^2 + bx + c$ has a repeated non-real root.

Observation C.3.24

Consider the homogeneous second order constant coefficient ODE

$$ay'' + by' + cy = 0$$

given by the differential operator $aD^2 + bD + cI$. Let r be a (possibly non-real) solution to $ax^2 + bx + c = 0$:

- e^{rt} is a particular solution of the ODE.
- If r is a double root, te^{rt} is also a particular solution.
- if $r = \alpha + \beta i$ is not real, Euler's formula allows us to express the real-valued solutions in terms of $\sin(\beta t)$ and $\cos(\beta t)$.

Due to the usefulness of its solutions, $ax^2 + bx + c = 0$ is called the **auxiliary equation** for this ODE.

Module C Section 4

Module C

Section C.1

Section C.2

Section C.3

Section C.4

Section C.5

Section C.6

Remark C.4.1

While first or second-order constant-coefficient ODEs usually solve to general solutions such as $y = c_1 e^t + c_2 e^{-2t}$, the values of the parameters c_1, c_2 may be determined when given additional information.

Module C

Section C.1

Section C.2

Section C.3

Section C.4

Section C.5

Section C.6

Activity C.4.2 (*~10 min*)

Solve the IVP

$$y' + 3y = 0, \quad y(0) = 2.$$

Module C

Section C.1

Section C.2

Section C.3

Section C.4

Section C.5

Section C.6

Activity C.4.3 (*~15 min*)

Solve $y'' - 6y' + 9y = 0$ where $y(0) = 2$ and $y(1) = \frac{3}{e^3}$.

Module C

Section C.1

Section C.2

Section C.3

Section C.4

Section C.5

Section C.6

Activity C.4.4 (*~15 min*)

Solve $y'' - 6y' + 8y = 0$ where $y(0) = 1$ and $y'(0) = -2$.

Module C

Section C.1

Section C.2

Section C.3

Section C.4

Section C.5

Section C.6

Module C Section 5

Observation C.5.1

Consider the homogeneous second order constant coefficient ODE

$$ay'' + by' + cy = 0.$$

- If r is a root of $ax^2 + bx + c = 0$, then e^{rt} is a solution of the ODE.
- If r is a double root (that is, $ax^2 + bx + c = (x - r)^2$), te^{rt} is also a solution.
- If $r = a + bi$ is not real, Euler's formula allows us to express e^{at+bit} in terms of e^{at} , $\sin(bt)$, and $\cos(bt)$ to get a real-valued general solution.

Activity C.5.2 (*~15 min*)

Consider the following scenario: a mass of 4 kg suspended from a damped spring with spring constant $k = 2 \text{ kg/s}^2$ and damping constant $b = 6 \text{ kg/s}$. As previously discussed, this is modeled by the ODE

$$my'' = -by' - ky.$$

Activity C.5.2 (*~15 min*)

Consider the following scenario: a mass of 4 kg suspended from a damped spring with spring constant $k = 2 \text{ kg/s}^2$ and damping constant $b = 6 \text{ kg/s}$. As previously discussed, this is modeled by the ODE

$$my'' = -by' - ky.$$

Part 1: Find the general solution for the ODE in terms of m, b, k .

Activity C.5.2 (*~15 min*)

Consider the following scenario: a mass of 4 kg suspended from a damped spring with spring constant $k = 2 \text{ kg/s}^2$ and damping constant $b = 6 \text{ kg/s}$. As previously discussed, this is modeled by the ODE

$$my'' = -by' - ky.$$

Part 1: Find the general solution for the ODE in terms of m, b, k .

Part 2: The mass is pulled down 0.3 m from its natural length and released from rest. Use the initial conditions $y(0) = ?$ and $y'(0) = ?$ to find the particular solution modeling this scenario.

Activity C.5.3 (*~5 min*)

A 1 kg mass is suspended from a spring with spring constant $k = 9 \text{ kg/s}^2$. No damping is applied, but an external electromagnetic force of $F(t) = \sin(t)$ is applied. Which of these ODEs models this scenario?

a) $my'' + ky + \sin(t) = 0$

b) $my'' + ky = \sin(t)$

c) $my'' + by' = \sin(t)$

d) $my'' + by' + \sin(t) = 0$

Observation C.5.4

Because my'' is the total force acting on the object, $-by' - ky$ is the force acting on the object by the spring, and an additional external force of $F(t)$ is applied, we get $my'' = -by' - ky + F(t)$ which rearranges to

$$my'' + ky = \sin(t)$$

when $b = 0$ (no damping) and $F(t) = \sin(t)$.

This is an example of a **nonhomogeneous** second-order constant coefficient equation of the form

$$ay'' + by' + cy = F(t)$$

since the $F(t) = \sin(t)$ term is not a multiple of y or its derivatives. As with first-order examples, these may be solved with variation of parameters.

Activity C.5.5 (*~15 min*)

Suppose y_1 and y_2 are two independent particular solutions of $ay'' + by' + cy = 0$.

By variation of parameters, we'll assume we can find a particular solution $y_p = v_1y_1 + v_2y_2$ for the ODE using the currently unknown functions v_1, v_2 .

Activity C.5.5 (*~15 min*)

Suppose y_1 and y_2 are two independent particular solutions of $ay'' + by' + cy = 0$.

By variation of parameters, we'll assume we can find a particular solution $y_p = v_1y_1 + v_2y_2$ for the ODE using the currently unknown functions v_1, v_2 .

Part 1: Use the product rule (on v_1y_1 and v_2y_2) to compute y_p' .

Activity C.5.5 (*~15 min*)

Suppose y_1 and y_2 are two independent particular solutions of $ay'' + by' + cy = 0$.

By variation of parameters, we'll assume we can find a particular solution $y_p = v_1y_1 + v_2y_2$ for the ODE using the currently unknown functions v_1, v_2 .

Part 1: Use the product rule (on v_1y_1 and v_2y_2) to compute y_p' .

Part 2: Since we get to choose what v_1, v_2 are, let's only look for examples where $v_1'y_1 + v_2'y_2 = 0$ to simplify calculations. Assuming this, compute y_p'' .

Activity C.5.5 (*~15 min*)

Suppose y_1 and y_2 are two independent particular solutions of $ay'' + by' + cy = 0$.

By variation of parameters, we'll assume we can find a particular solution $y_p = v_1y_1 + v_2y_2$ for the ODE using the currently unknown functions v_1, v_2 .

Part 1: Use the product rule (on v_1y_1 and v_2y_2) to compute y'_p .

Part 2: Since we get to choose what v_1, v_2 are, let's only look for examples where $v'_1y_1 + v'_2y_2 = 0$ to simplify calculations. Assuming this, compute y''_p .

Part 3: Simplify the ODE $ay''_p + by'_p + cy_p = f(x)$, keeping in mind that $ay''_1 + by'_1 + cy_1 = 0$ and $ay''_2 + by'_2 + cy_2 = 0$.

Observation C.5.6

If we can find functions v_1 and v_2 that solve the system of equations

$$y_1 v_1' + y_2 v_2' = 0$$

$$y_1' v_1 + y_2' v_2 = \frac{1}{a} f(t)$$

then $y_p = y_1 v_1 + y_2 v_2$ is a particular solution for $ay'' + by' + cy = f(x)$.

Activity C.5.7 (*~20 min*)

Consider the nonhomogeneous ODE

$$y'' + 9y = \sin(t)$$

of the form $ay'' + by' + cy = f(t)$ for $a = 1, b = 0, c = 9, f(t) = \sin(t)$.

Activity C.5.7 (~ 20 min)

Consider the nonhomogeneous ODE

$$y'' + 9y = \sin(t)$$

of the form $ay'' + by' + cy = f(t)$ for $a = 1, b = 0, c = 9, f(t) = \sin(t)$.

Part 1: Find $y_h = k_1y_1 + k_2y_2$, where y_1, y_2 are independent real-valued particular solutions of $y_h'' + 9y_h = 0$.

Activity C.5.7 (~ 20 min)

Consider the nonhomogeneous ODE

$$y'' + 9y = \sin(t)$$

of the form $ay'' + by' + cy = f(t)$ for $a = 1$, $b = 0$, $c = 9$, $f(t) = \sin(t)$.

Part 1: Find $y_h = k_1y_1 + k_2y_2$, where y_1, y_2 are independent real-valued particular solutions of $y_h'' + 9y_h = 0$.

Part 2: Substitute $a, f(t), y_1, y_2, y_1', y_2'$ into

$$y_1 v_1' + y_2 v_2' = 0$$

$$y_1' v_1' + y_2' v_2' = \frac{1}{a} f(t)$$

Activity C.5.7 (~ 20 min)

Consider the nonhomogeneous ODE

$$y'' + 9y = \sin(t)$$

of the form $ay'' + by' + cy = f(t)$ for $a = 1$, $b = 0$, $c = 9$, $f(t) = \sin(t)$.

Part 1: Find $y_h = k_1y_1 + k_2y_2$, where y_1, y_2 are independent real-valued particular solutions of $y_h'' + 9y_h = 0$.

Part 2: Substitute $a, f(t), y_1, y_2, y_1', y_2'$ into

$$\begin{aligned} y_1 v_1' + y_2 v_2' &= 0 \\ y_1' v_1 + y_2' v_2 &= \frac{1}{a} f(t) \end{aligned}$$

Part 3: Find v_1, v_2 by solving that system, and using

$$\begin{aligned} \int \sin(t) \cos(3t) dt &= \frac{1}{8} \cos(t) \cos(3t) + \frac{3}{8} \sin(t) \sin(3t) + C \text{ and} \\ \int \sin(t) \sin(3t) dt &= -\frac{1}{8} \cos(t) \sin(3t) + \frac{3}{8} \sin(t) \cos(3t) + C. \end{aligned}$$

Activity C.5.7 (~20 min)

Consider the nonhomogeneous ODE

$$y'' + 9y = \sin(t)$$

of the form $ay'' + by' + cy = f(t)$ for $a = 1, b = 0, c = 9, f(t) = \sin(t)$.

Part 1: Find $y_h = k_1y_1 + k_2y_2$, where y_1, y_2 are independent real-valued particular solutions of $y_h'' + 9y_h = 0$.

Part 2: Substitute $a, f(t), y_1, y_2, y_1', y_2'$ into

$$\begin{aligned} y_1 v_1' + y_2 v_2' &= 0 \\ y_1' v_1 + y_2' v_2 &= \frac{1}{a} f(t) \end{aligned}$$

Part 3: Find v_1, v_2 by solving that system, and using

$$\int \sin(t) \cos(3t) dt = \frac{1}{8} \cos(t) \cos(3t) + \frac{3}{8} \sin(t) \sin(3t) + C \text{ and}$$

$$\int \sin(t) \sin(3t) dt = -\frac{1}{8} \cos(t) \sin(3t) + \frac{3}{8} \sin(t) \cos(3t) + C.$$

Part 4: Use $y_p = y_1 v_1 + y_2 v_2$ to write the general solution $y = y_h + y_p$ of the original nonhomogeneous ODE.

Activity C.5.8 (*~10 min*)

Consider the nonhomogeneous ODE $y'' + 9y = \sin(3t)$.

Activity C.5.8 (~ 10 min)

Consider the nonhomogeneous ODE $y'' + 9y = \sin(3t)$.

Part 1: Find v_1 and v_2 by solving

$$y_1 v_1' + y_2 v_2' = 0$$

$$y_1' v_1 + y_2' v_2 = \frac{1}{a} f(t)$$

for particular solutions y_1, y_2 of $y_h'' + 9y_h = 0$. Use

$$\int \sin(3t) \cos(3t) dt = \frac{1}{6} \sin^2(3t) + C \text{ and}$$

$$\int \sin^2(3t) dt = \frac{1}{6} (3t - \sin(3t) \cos(3t)) + C.$$

Activity C.5.8 (~ 10 min)

Consider the nonhomogeneous ODE $y'' + 9y = \sin(3t)$.

Part 1: Find v_1 and v_2 by solving

$$y_1 v_1' + y_2 v_2' = 0$$

$$y_1' v_1' + y_2' v_2' = \frac{1}{a} f(t)$$

for particular solutions y_1, y_2 of $y_h'' + 9y_h = 0$. Use

$$\int \sin(3t) \cos(3t) dt = \frac{1}{6} \sin^2(3t) + C \text{ and}$$

$$\int \sin^2(3t) dt = \frac{1}{6} (3t - \sin(3t) \cos(3t)) + C.$$

Part 2: Write the general solution of the original nonhomogeneous ODE.

Module C Section 6

Activity C.6.1 (*~20 min*)

A 3kg mass is attached to a spring requires 4 Newtons ($\text{kg} \cdot \text{m}/\text{s}^2$) to pull the mass 2 meters from its natural length. No damper is applied. The mass is then released from rest.

Activity C.6.1 (*~20 min*)

A 3kg mass is attached to a spring requires 4 Newtons ($\text{kg} \cdot \text{m}/\text{s}^2$) to pull the mass 2 meters from its natural length. No damper is applied. The mass is then released from rest.

Part 1: Adapt the ODE

$$my'' + by' + ky = 0$$

to give an initial value problem modeling this scenario.

Activity C.6.1 (*~20 min*)

A 3kg mass is attached to a spring requires 4 Newtons ($\text{kg} \cdot \text{m}/\text{s}^2$) to pull the mass 2 meters from its natural length. No damper is applied. The mass is then released from rest.

Part 1: Adapt the ODE

$$my'' + by' + ky = 0$$

to give an initial value problem modeling this scenario.

Part 2: How much time will pass before the spring first returns to its natural length?