

Sample Assessment Exercises

This document contains one exercise and solution for each standard. The goal is to give you an idea of what the exercises might look like, and what the expectations for a complete solution are.

E1. Write a system of linear equations corresponding to the following augmented matrix.

$$\left[\begin{array}{cccc|c} 3 & 2 & 0 & 1 & 1 \\ -1 & -4 & 1 & -7 & 0 \\ 0 & 1 & -1 & 0 & -2 \end{array} \right]$$

Solution:

$$\begin{aligned} 3x_1 + 2x_2 &+ x_4 = 1 \\ -x_1 - 4x_2 + x_3 - 7x_4 &= 0 \\ x_2 - x_3 &= -2 \end{aligned}$$

□

E2. Put the following matrix in reduced row echelon form.

$$\begin{bmatrix} 0 & 3 & 1 & 2 \\ 1 & 2 & -1 & -3 \\ 2 & 4 & -1 & -1 \end{bmatrix}$$

Solution:

$$\begin{aligned} \begin{bmatrix} 0 & 3 & 1 & 2 \\ 1 & 2 & -1 & -3 \\ 2 & 4 & -1 & -1 \end{bmatrix} &\sim \begin{bmatrix} \textcircled{1} & 2 & -1 & -3 \\ 0 & 3 & 1 & 2 \\ 2 & 4 & -1 & -1 \end{bmatrix} && \text{Swap Rows 1 and 2} \\ &\sim \begin{bmatrix} \textcircled{1} & 2 & -1 & -3 \\ 0 & 3 & 1 & 2 \\ 0 & 0 & 1 & 5 \end{bmatrix} && \text{Add } -2 \text{ Row 1 to Row 3} \\ &\sim \begin{bmatrix} \textcircled{1} & 2 & -1 & -3 \\ 0 & \textcircled{1} & \frac{1}{3} & \frac{2}{3} \\ 0 & 0 & 1 & 5 \end{bmatrix} && \text{Multiply Row 3 by } \frac{1}{3} \\ &\sim \begin{bmatrix} \textcircled{1} & 0 & -\frac{5}{3} & -\frac{13}{3} \\ 0 & \textcircled{1} & \frac{1}{3} & \frac{2}{3} \\ 0 & 0 & \textcircled{1} & 5 \end{bmatrix} && \text{Add } -2 \text{ Row 2 to Row 1} \\ &\sim \begin{bmatrix} \textcircled{1} & 0 & -\frac{5}{3} & -\frac{13}{3} \\ 0 & \textcircled{1} & 0 & -1 \\ 0 & 0 & \textcircled{1} & 5 \end{bmatrix} && \text{Add } -\frac{1}{3} \text{ Row 3 to Row 2} \\ &\sim \begin{bmatrix} \textcircled{1} & 0 & 0 & 4 \\ 0 & \textcircled{1} & 0 & -1 \\ 0 & 0 & \textcircled{1} & 5 \end{bmatrix} && \text{Add } \frac{5}{3} \text{ Row 3 to Row 1} \end{aligned}$$

□

E3. Show how to find the solution set for the following system of linear equations.

$$\begin{aligned} 2x + 4y + z &= 5 \\ x + 2y &= 3 \end{aligned}$$

Solution: First, note that this system corresponds to the matrix $\left(\left[\begin{array}{ccc|c} 2 & 4 & 1 & 5 \\ 1 & 2 & 0 & 3 \end{array}\right]\right)$. Then we compute (using technology)

$$\text{RREF}\left(\left[\begin{array}{ccc|c} 2 & 4 & 1 & 5 \\ 1 & 2 & 0 & 3 \end{array}\right]\right) = \left[\begin{array}{ccc|c} 1 & 2 & 0 & 3 \\ 0 & 0 & 1 & -1 \end{array}\right].$$

This corresponds to the system

$$\begin{aligned} x + 2y &= 3 \\ z &= -1 \end{aligned}$$

Since the y -column is a non-pivot column, it is a free variable, so we let $y = a$; then we have

$$\begin{aligned} x + 2y &= 3 \\ y &= a \\ z &= -1 \end{aligned}$$

and thus

$$\begin{aligned} x &= 3 - 2a \\ y &= a \\ z &= -1 \end{aligned}$$

So the solution set is

$$\left\{ \left[\begin{array}{c} 3 - 2a \\ a \\ -1 \end{array} \right] \mid a \in \mathbb{R} \right\}$$

□

V1. Let V be the set of all pairs of numbers (x, y) of real numbers together with the following operations:

$$\begin{aligned} (x_1, y_2) \oplus (x_2, y_2) &= (2x_1 + 2x_2, 2y_1 + 2y_2) \\ c \odot (x, y) &= (cx, c^2y) \end{aligned}$$

(a) Show that scalar multiplication distributes over vector addition:

$$c \odot ((x_1, y_1) \oplus (x_2, y_2)) = c \odot (x_1, y_1) \oplus c \odot (x_2, y_2)$$

(b) Explain why V nonetheless is not a vector space.

Solution:

(a) We compute both sides:

$$\begin{aligned}c \odot ((x_1, y_1) \oplus (x_2, y_2)) &= c \odot (2x_1 + 2x_2, 2y_1 + 2y_2) \\ &= (c(2x_1 + 2x_2), c^2(2y_1 + 2y_2)) \\ &= (2cx_1 + 2cx_2, 2c^2y_1 + 2c^2y_2)\end{aligned}$$

and

$$\begin{aligned}c \odot (x_1, y_1) \oplus c \odot (x_2, y_2) &= (cx_1, c^2y_1) \oplus (cx_2, c^2y_2) \\ &= (2cx_1 + 2cx_2, 2c^2y_1 + 2c^2y_2)\end{aligned}$$

Since these are the same, we have shown that the property holds.

(b) To show V is not a vector space, we must show that it fails one of the 8 defining properties of vector spaces. We will show that scalar multiplication does not distribute over scalar addition, i.e., there are values such that

$$(c + d) \odot (x, y) \neq c \odot (x, y) \oplus d \odot (x, y)$$

(*Solution method 1*) First, we compute

$$\begin{aligned}(c + d) \odot (x, y) &= ((c + d)x, (c + d)^2y) \\ &= ((c + d)x, (c^2 + 2cd + d^2)y).\end{aligned}$$

Then we compute

$$\begin{aligned}c \odot (x, y) \oplus d \odot (x, y) &= (cx, c^2y) \oplus (dx, d^2y) \\ &= (2cx + 2dx, 2c^2y + 2d^2y).\end{aligned}$$

Since $(c + d)x \neq 2cx + 2dx$ when $c, d, x, y = 1$, the property fails to hold.

(*Solution method 2*) When we let $c, d, x, y = 1$, we may simplify both sides as follows.

$$\begin{aligned}(c + d) \odot (x, y) &= 2 \odot (1, 1) \\ &= (2 \cdot 1, 2^2 \cdot 1) \\ &= (2, 4)\end{aligned}$$

$$\begin{aligned}c \odot (x, y) \oplus d \odot (x, y) &= 1 \odot (1, 1) \oplus 1 \odot (1, 1) \\ &= (1 \cdot 1, 1^2 \cdot 1) \oplus (1 \cdot 1, 1^2 \cdot 1) \\ &= (1, 1) \oplus (1, 1) \\ &= (2 \cdot 1 + 2 \cdot 1, 2 \cdot 1 + 2 \cdot 1) \\ &= (4, 4)\end{aligned}$$

Since these ordered pairs are different, the property fails to hold.

□

V2. Explain why the vector $\begin{bmatrix} 3 \\ -1 \\ 2 \end{bmatrix}$ is or is not a linear combination of the vectors $\begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}$, $\begin{bmatrix} 3 \\ 2 \\ -1 \end{bmatrix}$, and $\begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix}$.

Solution: By definition, the statement

$$\begin{bmatrix} 3 \\ -1 \\ 2 \end{bmatrix} \text{ is a linear combination of the vectors } \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 3 \\ 2 \\ -1 \end{bmatrix}, \text{ and } \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix}$$

is equivalent to the statement

$$\text{There exists a solution to the system of equations } x_1 \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} + x_2 \begin{bmatrix} 3 \\ 2 \\ -1 \end{bmatrix} + x_3 \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix} = \begin{bmatrix} 3 \\ -1 \\ 2 \end{bmatrix}.$$

This system corresponds to the augmented matrix $\left[\begin{array}{ccc|c} 1 & 3 & 1 & 3 \\ 0 & 2 & 1 & -1 \\ 1 & -1 & -1 & 2 \end{array} \right]$. Therefore, we compute

$$\text{RREF} \left[\begin{array}{ccc|c} 1 & 3 & 1 & 3 \\ 0 & 2 & 1 & -1 \\ 1 & -1 & -1 & 2 \end{array} \right] = \left[\begin{array}{ccc|c} 1 & 0 & -\frac{1}{2} & 0 \\ 0 & 1 & \frac{1}{2} & 0 \\ 0 & 0 & 0 & 1 \end{array} \right].$$

Since this corresponds to an inconsistent system of equations, the system of equations

$$x_1 \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} + x_2 \begin{bmatrix} 3 \\ 2 \\ -1 \end{bmatrix} + x_3 \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix} = \begin{bmatrix} 3 \\ -1 \\ 2 \end{bmatrix}$$

has no solution, and therefore $\begin{bmatrix} 3 \\ -1 \\ 2 \end{bmatrix}$ is not a linear combination of $\begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}$, $\begin{bmatrix} 3 \\ 2 \\ -1 \end{bmatrix}$, and $\begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix}$.

□

V3. Explain why the vectors $\begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}$, $\begin{bmatrix} 3 \\ 2 \\ -1 \end{bmatrix}$, and $\begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix}$ span or don't span \mathbb{R}^3 .

Solution: By definition, the statement

$$\text{The vectors } \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 3 \\ 2 \\ -1 \end{bmatrix}, \text{ and } \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix} \text{ span } \mathbb{R}^3$$

is equivalent to the statement

$$\text{For every } \vec{v} \in \mathbb{R}^3, \text{ the system } x_1 \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} + x_2 \begin{bmatrix} 3 \\ 2 \\ -1 \end{bmatrix} + x_3 \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix} = \vec{v} \text{ has a solution.}$$

We compute

$$\text{RREF} \left[\begin{array}{ccc} 1 & 3 & 1 \\ 0 & 2 & 1 \\ 1 & -1 & -1 \end{array} \right] = \left[\begin{array}{ccc} 1 & 0 & -\frac{1}{2} \\ 0 & 1 & \frac{1}{2} \\ 0 & 0 & 0 \end{array} \right]$$

Since the last row lacks a pivot, there is some vector $\vec{v} \in \mathbb{R}^3$ that upon augmenting this matrix will produce an inconsistent system. That vector will not be in the span of these three vectors, so the vectors do not span \mathbb{R}^3 .

□

V4. Consider the following two sets of Euclidean vectors.

$$W = \left\{ \begin{bmatrix} x \\ y \\ z \\ w \end{bmatrix} \mid x + y = 3z + 2w \right\} \quad U = \left\{ \begin{bmatrix} x \\ y \\ z \\ w \end{bmatrix} \mid x + y = 3z + w^2 \right\}$$

Explain why one of these sets is a subspace of \mathbb{R}^3 , and why the other is not.

Solution:

To show that W is a subspace, let $\vec{v} = \begin{bmatrix} x_1 \\ y_1 \\ z_1 \\ w_1 \end{bmatrix} \in W$ and $\vec{w} = \begin{bmatrix} x_2 \\ y_2 \\ z_2 \\ w_2 \end{bmatrix} \in W$, so we know that $x_1 + y_1 = 3z_1 + 2w_1$ and $x_2 + y_2 = 3z_2 + 2w_2$. Consider

$$\begin{bmatrix} x_1 \\ y_1 \\ z_1 \\ w_1 \end{bmatrix} + \begin{bmatrix} x_2 \\ y_2 \\ z_2 \\ w_2 \end{bmatrix} = \begin{bmatrix} x_1 + x_2 \\ y_1 + y_2 \\ z_1 + z_2 \\ w_1 + w_2 \end{bmatrix}.$$

To see if $\vec{v} + \vec{w} \in W$, we need to check if $(x_1 + x_2) + (y_1 + y_2) = 3(z_1 + z_2) + 2(w_1 + w_2)$. We compute

$$\begin{aligned} (x_1 + x_2) + (y_1 + y_2) &= (x_1 + y_1) + (x_2 + y_2) && \text{by regrouping} \\ &= (3z_1 + 2w_1) + (3z_2 + 2w_2) && \text{since } \vec{v}, \vec{w} \in W \\ &= 3(z_1 + z_2) + 2(w_1 + w_2) && \text{by regrouping.} \end{aligned}$$

Thus $\vec{v} + \vec{w} \in W$, so W is closed under vector addition.

Now consider

$$c\vec{v} = \begin{bmatrix} cx_1 \\ cy_1 \\ cz_1 \\ cw_1 \end{bmatrix}.$$

Similarly, to check that $c\vec{v} \in W$, we need to check if $cx_1 + cy_1 = 3(cz_1) + 2(cw_1)$, so we compute

$$\begin{aligned} cx_1 + cy_1 &= c(x_1 + y_1) && \text{by factoring} \\ &= c(3z_1 + 2w_1) && \text{since } \vec{v} \in W \\ &= 3(cz_1) + 2(cw_1) && \text{by regrouping} \end{aligned}$$

and we see that $c\vec{v} \in W$, so W is closed under scalar multiplication. Therefore W is a subspace of \mathbb{R}^3 .

Now, to show U is not a subspace, we will show that it is not closed under vector addition.

(Solution Method 1) Now let $\vec{v} = \begin{bmatrix} x_1 \\ y_1 \\ z_1 \\ w_1 \end{bmatrix} \in U$ and $\vec{w} = \begin{bmatrix} x_2 \\ y_2 \\ z_2 \\ w_2 \end{bmatrix} \in U$, so we know that $x_1 + y_1 = 3z_1 + w_1^2$ and $x_2 + y_2 = 3z_2 + w_2^2$. Consider

$$\vec{v} + \vec{w} = \begin{bmatrix} x_1 \\ y_1 \\ z_1 \\ w_1 \end{bmatrix} + \begin{bmatrix} x_2 \\ y_2 \\ z_2 \\ w_2 \end{bmatrix} = \begin{bmatrix} x_1 + x_2 \\ y_1 + y_2 \\ z_1 + z_2 \\ w_1 + w_2 \end{bmatrix}.$$

To see if $\vec{v} + \vec{w} \in U$, we need to check if $(x_1 + x_2) + (y_1 + y_2) = 3(z_1 + z_2) + (w_1 + w_2)^2$. We compute

$$\begin{aligned} (x_1 + x_2) + (y_1 + y_2) &= (x_1 + y_1) + (x_2 + y_2) && \text{by regrouping} \\ &= (3z_1 + w_1^2) + (3z_2 + w_2^2) && \text{since } \vec{v}, \vec{w} \in W \\ &= 3(z_1 + z_2) + (w_1^2 + w_2^2) && \text{by regrouping} \end{aligned}$$

and thus $\vec{v} + \vec{w} \in U$ **only when** $w_1^2 + w_2^2 = (w_1 + w_2)^2$. Since this is not true in general, U is not closed under vector addition, and thus cannot be a subspace.

(*Solution Method 2*) Note that the vector $\vec{v} = \begin{bmatrix} 0 \\ 1 \\ 0 \\ 1 \end{bmatrix}$ belongs to U since $0 + 1 = 3(0) + 1^2$. However, the vector

$2\vec{v} = \begin{bmatrix} 0 \\ 2 \\ 0 \\ 2 \end{bmatrix}$ does not belong to U since $0 + 2 \neq 3(0) + 2^2$. Therefore U is not closed under scalar multiplication, and thus is not a subspace. □

V5. Explain why the set of vectors

$$\left\{ \begin{bmatrix} 3 \\ 2 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} -1 \\ 1 \\ 2 \\ 3 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ -1 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 5 \\ 1 \\ 5 \end{bmatrix} \right\}$$

is linearly dependent or linearly independent.

Solution: The vectors $\begin{bmatrix} 3 \\ 2 \\ 1 \\ 0 \end{bmatrix}$, $\begin{bmatrix} -1 \\ 1 \\ 2 \\ 3 \end{bmatrix}$, $\begin{bmatrix} 0 \\ 1 \\ -1 \\ 1 \end{bmatrix}$, and $\begin{bmatrix} 2 \\ 5 \\ 1 \\ 5 \end{bmatrix}$ are linearly independent precisely when the system of equations

$$x_1 \begin{bmatrix} 3 \\ 2 \\ 1 \\ 0 \end{bmatrix} + x_2 \begin{bmatrix} -1 \\ 1 \\ 2 \\ 3 \end{bmatrix} + x_3 \begin{bmatrix} 0 \\ 1 \\ -1 \\ 1 \end{bmatrix} + x_4 \begin{bmatrix} 2 \\ 5 \\ 1 \\ 5 \end{bmatrix} = \vec{0}$$

has a unique solution (namely, $\vec{0}$).

Converting the left side of this system to the corresponding matrix and row reducing, we have

$$\text{RREF} \begin{bmatrix} 3 & -1 & 0 & 2 \\ 2 & 1 & 1 & 5 \\ 1 & 2 & -1 & 1 \\ 0 & 3 & 1 & 5 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

Since the fourth column is not a pivot column, the system has (infinitely many) nontrivial solutions. Thus the set of vectors is linearly dependent. □

V6. Explain why the set of vectors

$$\left\{ \begin{bmatrix} 3 \\ 2 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} -1 \\ 1 \\ 2 \\ 3 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ -1 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 5 \\ 1 \\ 5 \end{bmatrix} \right\}$$

is or is not a basis of \mathbb{R}^4 .

Solution: Compute

$$\text{RREF} \begin{bmatrix} 3 & -1 & 0 & 2 \\ 2 & 1 & 1 & 5 \\ 1 & 2 & -1 & 1 \\ 0 & 3 & 1 & 5 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

Since the fourth column is not a pivot column, the vectors are linearly dependent and thus not a basis of \mathbb{R}^4 .

(Alternate solution:) Since the fourth row is not a pivot row, the vectors do not span \mathbb{R}^4 and thus are not a basis of \mathbb{R}^4 .

□

V7. Find a basis for the subspace

$$W = \text{span} \left\{ \begin{bmatrix} 1 \\ -3 \\ -1 \\ 2 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 1 \\ -2 \end{bmatrix}, \begin{bmatrix} 3 \\ -6 \\ -1 \\ 2 \end{bmatrix}, \begin{bmatrix} 1 \\ 6 \\ 1 \\ -1 \end{bmatrix}, \begin{bmatrix} 2 \\ 3 \\ 0 \\ 1 \end{bmatrix} \right\}.$$

Be sure to explain why your subspace is a basis.

Solution: Observe that

$$\text{RREF} \begin{bmatrix} 1 & 1 & 3 & 1 & 2 \\ -3 & 0 & -6 & 6 & 3 \\ -1 & 1 & -1 & 1 & 0 \\ 2 & -2 & 2 & -1 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 2 & 0 & 1 \\ 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

If we remove the vectors yielding non-pivot columns, the resulting set will span the same vectors while being linearly independent. Therefore

$$\left\{ \begin{bmatrix} 1 \\ -3 \\ -1 \\ 2 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 1 \\ -2 \end{bmatrix}, \begin{bmatrix} 1 \\ 6 \\ 1 \\ -1 \end{bmatrix} \right\}$$

is a basis of W .

□

V8. Explain how to find the dimension of

$$W = \text{span} \left\{ \begin{bmatrix} 1 \\ -3 \\ -1 \\ 2 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 1 \\ -2 \end{bmatrix}, \begin{bmatrix} 3 \\ -6 \\ -1 \\ 2 \end{bmatrix}, \begin{bmatrix} 1 \\ 6 \\ 1 \\ -1 \end{bmatrix}, \begin{bmatrix} 2 \\ 3 \\ 0 \\ 1 \end{bmatrix} \right\}.$$

Solution: The dimension of a space is equal to how many vectors belong to any basis for the space. So since

$$\text{RREF} \begin{bmatrix} 1 & 1 & 3 & 1 & 2 \\ -3 & 0 & -6 & 6 & 3 \\ -1 & 1 & -1 & 1 & 0 \\ 2 & -2 & 2 & -1 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 2 & 0 & 1 \\ 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

has three pivot columns, any basis of W has three elements, and therefore $\dim W = 3$.

□

V9. Find a basis for the subspace of \mathcal{P}^3

$$W = \text{span} \{3x^3 + 2x^2 + x, -x^3 + x^2 + 2x + 3, x^2 - x + 1, 2x^3 + 5x^2 + x + 5\}.$$

Be sure to explain why your result is a basis.

Solution: This question is equivalent to finding a basis for the subspace

$$W' = \text{span} \left\{ \begin{bmatrix} 3 \\ 2 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} -1 \\ 1 \\ 2 \\ 3 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ -1 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 5 \\ 1 \\ 5 \end{bmatrix} \right\}$$

of Euclidean vectors.

Compute

$$\text{RREF} \begin{bmatrix} 3 & -1 & 0 & 2 \\ 2 & 1 & 1 & 5 \\ 1 & 2 & -1 & 1 \\ 0 & 3 & 1 & 5 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

Since the fourth column is not a pivot column, a basis for W' is given by

$$\left\{ \begin{bmatrix} 3 \\ 2 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} -1 \\ 1 \\ 2 \\ 3 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ -1 \\ 1 \end{bmatrix} \right\}$$

Thus a basis for W is given by

$$\{3x^3 + 2x^2 + x, -x^3 + x^2 + 2x + 3, x^2 - x + 1\}$$

□

V10. Explain how to find a basis for the solution set of the homogeneous system of equations

$$\begin{aligned} x_1 + x_2 + 3x_3 + x_4 + 2x_5 &= 0 \\ -3x_1 - 6x_3 + 6x_4 + 3x_5 &= 0 \\ -x_1 + x_2 - x_3 + x_4 &= 0 \\ 2x_1 - 2x_2 + 2x_3 - x_4 + x_5 &= 0. \end{aligned}$$

Solution: Observe that

$$\text{RREF} \left[\begin{array}{ccccc|c} 1 & 1 & 3 & 1 & 2 & 0 \\ -3 & 0 & -6 & 6 & 3 & 0 \\ -1 & 1 & -1 & 1 & 0 & 0 \\ 2 & -2 & 2 & -1 & 1 & 0 \end{array} \right] = \left[\begin{array}{ccccc|c} 1 & 0 & 2 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{array} \right]$$

Letting $x_3 = a$ and $x_5 = b$ (since those correspond to the non-pivot columns), this is equivalent to the system

$$\begin{aligned} x_1 + 2x_3 + x_5 &= 0 \\ x_2 + x_3 &= 0 \\ x_3 &= a \\ x_4 + x_5 &= 0 \\ x_5 &= b \end{aligned}$$

Thus, the solution set is

$$\left\{ \left[\begin{array}{c} -2a - b \\ -a \\ a \\ -b \\ b \end{array} \right] \middle| a, b \in \mathbb{R} \right\}.$$

Since we can write

$$\left[\begin{array}{c} -2a - b \\ -a \\ a \\ -b \\ b \end{array} \right] = a \left[\begin{array}{c} -2 \\ -1 \\ 1 \\ 0 \\ 0 \end{array} \right] + b \left[\begin{array}{c} -1 \\ 0 \\ 0 \\ -1 \\ 1 \end{array} \right],$$

a basis for the solution space is

$$\left\{ \left[\begin{array}{c} -2 \\ -1 \\ 1 \\ 0 \\ 0 \end{array} \right], \left[\begin{array}{c} -1 \\ 0 \\ 0 \\ -1 \\ 1 \end{array} \right] \right\}.$$

□

A1. Consider the following maps of polynomials $S : \mathcal{P} \rightarrow \mathcal{P}$ and $T : \mathcal{P} \rightarrow \mathcal{P}$ defined by

$$S(f(x)) = 3xf(x) \text{ and } T(f(x)) = 3f'(x)f(x).$$

Explain why one of these maps is a linear transformation, and why the other map is not.

Solution: To show S is a linear transformation, we must show two things:

$$\begin{aligned} S(f(x) + g(x)) &= S(f(x)) + S(g(x)) \\ S(cf(x)) &= cS(f(x)) \end{aligned}$$

To show S respects addition, we compute

$$\begin{aligned} S(f(x) + g(x)) &= 3x(f(x) + g(x)) && \text{by definition of } S \\ &= 3xf(x) + 3xg(x) && \text{by distributing} \end{aligned}$$

But note that $S(f(x)) = 3xf(x)$ and $S(g(x)) = 3xg(x)$, so we have $S(f(x) + g(x)) = S(f(x)) + S(g(x))$.

For the second part, we compute

$$\begin{aligned} S(cf(x)) &= 3x(cf(x)) && \text{by definition of } S \\ &= 3cxf(x) && \text{rewriting the multiplication.} \end{aligned}$$

But note that $cS(f(x)) = c(3xf(x)) = 3cxf(x)$ as well, so we have $S(cf(x)) = cS(f(x))$. Now, since S respects both addition and scalar multiplication, we can conclude S is a linear transformation.

(*Solution method 1*) As for T , we compute

$$\begin{aligned} T(f(x) + g(x)) &= 3(f(x) + g(x))'(f(x) + g(x)) && \text{by definition of } T \\ &= 3(f'(x) + g'(x))(f(x) + g(x)) && \text{since the derivative is linear} \\ &= 3f(x)f'(x) + 3f(x)g'(x) + 3f'(x)g(x) + 3g(x)g'(x) && \text{by distributing} \end{aligned}$$

However, note that $T(f(x)) + T(g(x)) = 3f'(x)f(x) + 3g'(x)g(x)$, which is not always the same polynomial (for example, when $f(x) = g(x) = x$). So we see that $T(f(x) + g(x)) \neq T(f(x)) + T(g(x))$, so T does not respect addition and is therefore not a linear transformation.

(*Solution method 2*) As for T , we may choose the polynomial $f(x) = x$ and scalar $c = 2$. Then

$$T(cf(x)) = T(2x) = 3(2x)'(2x) = 3(2)(2x) = 12x.$$

But on the other hand,

$$cT(f(x)) = 2T(x) = 2(3)(2x)'(2x) = 2(3)(2)(2x) = 24x.$$

Since this isn't the same polynomial, T does not preserve multiplication and is therefore not a linear transformation.

□

A2. Let $T : \mathbb{R}^3 \rightarrow \mathbb{R}^4$ be the linear transformation given by

$$T\left(\begin{pmatrix} x \\ y \\ z \end{pmatrix}\right) = \begin{bmatrix} -x + y \\ -x + 3y - z \\ 7x + y + 3z \\ 0 \end{bmatrix}.$$

(a) Explain how to find the standard matrix for T .

(b) Explain how to find $T\left(\begin{pmatrix} -2 \\ 0 \\ 3 \end{pmatrix}\right)$

Solution:

(a) Since

$$T\left(\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}\right) = \begin{bmatrix} -1 \\ -1 \\ 7 \\ 0 \end{bmatrix} \quad T\left(\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}\right) = \begin{bmatrix} 1 \\ 3 \\ 1 \\ 0 \end{bmatrix} \quad T\left(\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}\right) = \begin{bmatrix} 0 \\ -1 \\ 3 \\ 0 \end{bmatrix}$$

The standard matrix is $\begin{bmatrix} -1 & 1 & 0 \\ -1 & 3 & -1 \\ 7 & 1 & 3 \\ 0 & 0 & 0 \end{bmatrix}$.

$$(b) T\left(\begin{bmatrix} -2 \\ 0 \\ 3 \end{bmatrix}\right) = \begin{bmatrix} -(-2) + (0) \\ -(-2) + 3(0) - (3) \\ 7(-2) + (0) + 3(3) \\ 0 \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \\ -5 \\ 0 \end{bmatrix}$$

Alternatively, $\begin{bmatrix} -1 & 1 & 0 \\ -1 & 3 & -1 \\ 7 & 1 & 3 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} -2 \\ 0 \\ 3 \end{bmatrix} = \begin{bmatrix} -1(-2) + 1(0) + 0(3) \\ -1(-2) + 3(0) - 1(3) \\ 7(-2) + 1(0) + 3(3) \\ 0(-2) + 0(0) + 0(3) \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \\ -5 \\ 0 \end{bmatrix}$.

□

A3. Let $T : \mathbb{R}^4 \rightarrow \mathbb{R}^3$ be the linear transformation given by

$$T\left(\begin{bmatrix} x \\ y \\ z \\ w \end{bmatrix}\right) = \begin{bmatrix} x + 3y + 2z - 3w \\ 2x + 4y + 6z - 10w \\ x + 6y - z + 3w \end{bmatrix}$$

Show how to compute a basis for the image and a basis for the kernel of T .

Solution: First, we note the standard matrix

$$A = \begin{bmatrix} 1 & 3 & 2 & -3 \\ 2 & 4 & 6 & -10 \\ 1 & 6 & -1 & 3 \end{bmatrix}$$

and compute

$$\text{RREF}(A) = \begin{bmatrix} 1 & 0 & 5 & -9 \\ 0 & 1 & -1 & 2 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

A basis for the image is given by the pivot columns, namely

$$\left\{ \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 3 \\ 4 \\ 6 \end{bmatrix} \right\}.$$

The kernel is the solution set of the corresponding homogeneous system of equations

$$\ker T = \left\{ \begin{bmatrix} -5a + 9b \\ a - 2b \\ a \\ b \end{bmatrix} \mid a, b \in \mathbb{R} \right\}$$

so a basis for the kernel is

$$\left\{ \begin{bmatrix} -5 \\ 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 9 \\ -2 \\ 0 \\ 1 \end{bmatrix} \right\}.$$

□

A4. Let $T : \mathbb{R}^4 \rightarrow \mathbb{R}^3$ be the linear transformation given by the standard matrix $\begin{bmatrix} 1 & 3 & 2 & -3 \\ 2 & 4 & 6 & -10 \\ 1 & 6 & -1 & 3 \end{bmatrix}$.

- (a) Explain why T is or is not injective.
- (b) Explain why T is or is not surjective.

Solution: Compute

$$\text{RREF} \begin{bmatrix} 1 & 3 & 2 & -3 \\ 2 & 4 & 6 & -10 \\ 1 & 6 & -1 & 3 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 5 & -9 \\ 0 & 1 & -1 & 2 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

- (a) Note that the third and fourth columns are non-pivot columns, which means $\ker T$ contains infinitely many vectors, so T is not injective.
- (b) Since the third row lacks a pivot, the image (i.e. the span of the columns) is a 2-dimensional subspace (and thus does not equal \mathbb{R}^3), so T is not surjective.

□

M1. Of the following three matrices, only two may be multiplied.

$$A = \begin{bmatrix} 1 & -3 \\ 0 & 1 \end{bmatrix} \qquad B = \begin{bmatrix} 4 & 1 & 2 \end{bmatrix} \qquad C = \begin{bmatrix} 0 & 1 & 3 \\ 1 & -2 & 5 \end{bmatrix}$$

Explain which two may be multiplied and why. Then show how to find their product.

Solution: AC is the only one that can be computed, since A is 2×2 and C is 2×3 . Thus AC will be the 2×3 matrix given by

$$\begin{aligned} AC(\vec{e}_1) &= A \begin{pmatrix} 0 \\ 1 \end{pmatrix} = 0 \begin{bmatrix} 1 \\ 0 \end{bmatrix} + 1 \begin{bmatrix} -3 \\ 1 \end{bmatrix} = \begin{bmatrix} -3 \\ 1 \end{bmatrix} \\ AC(\vec{e}_2) &= A \begin{pmatrix} 1 \\ -2 \end{pmatrix} = 1 \begin{bmatrix} 1 \\ 0 \end{bmatrix} - 2 \begin{bmatrix} -3 \\ 1 \end{bmatrix} = \begin{bmatrix} 7 \\ -2 \end{bmatrix} \\ AC(\vec{e}_3) &= A \begin{pmatrix} 3 \\ 5 \end{pmatrix} = 3 \begin{bmatrix} 1 \\ 0 \end{bmatrix} + 5 \begin{bmatrix} -3 \\ 1 \end{bmatrix} = \begin{bmatrix} -12 \\ 5 \end{bmatrix} \end{aligned}$$

Thus

$$AC = \begin{bmatrix} -3 & 7 & -12 \\ 1 & -2 & 5 \end{bmatrix}.$$

□

M2. Explain why the matrix $\begin{bmatrix} 1 & 3 & 2 \\ 2 & 4 & 6 \\ 1 & 6 & -1 \end{bmatrix}$ is or is not invertible.

Solution: We compute

$$\text{RREF} \left(\begin{bmatrix} 1 & 3 & 2 \\ 2 & 4 & 6 \\ 1 & 6 & -1 \end{bmatrix} \right) = \begin{bmatrix} 1 & 0 & 5 \\ 0 & 1 & -1 \\ 0 & 0 & 0 \end{bmatrix}.$$

Since its RREF is not the identity matrix, the linear map is not bijective and thus the matrix is not invertible.

□

M3. Show how to compute the inverse of the matrix $A = \begin{bmatrix} 1 & 2 & 3 & 5 \\ 0 & -1 & 4 & -2 \\ 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 1 \end{bmatrix}$.

Solution: To find the matrix A^{-1} where $AA^{-1} = I$, we need to solve the augmented matrix $[A|I]$.

$$\text{RREF} \left(\left[\begin{array}{cccc|cccc} 1 & 2 & 3 & 5 & 1 & 0 & 0 & 0 \\ 0 & -1 & 4 & -2 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 3 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 \end{array} \right] \right) = \left[\begin{array}{cccc|cccc} 1 & 0 & 0 & 0 & 1 & 2 & -11 & 32 \\ 0 & 1 & 0 & 0 & 0 & -1 & 4 & -14 \\ 0 & 0 & 1 & 0 & 0 & 0 & 1 & -3 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 \end{array} \right]$$

So the inverse is $\begin{bmatrix} 1 & 2 & -11 & 32 \\ 0 & -1 & 4 & -14 \\ 0 & 0 & 1 & -3 \\ 0 & 0 & 0 & 1 \end{bmatrix}$.

□

G1.

- Give a 3×3 matrix P that may be used to perform the row operation $R_3 \rightarrow R_3 + 4R_1$.
- Give a 3×3 matrix Q that may be used to perform the row operation $R_1 \rightarrow -4R_1$.
- Suppose A is a 3×3 matrix with determinant -5 . Use matrix multiplication to describe the matrix obtained by applying $R_3 \rightarrow 4R_1 + R_3$ and then $R_1 \rightarrow -4R_1$ to A (note the order).
- Finally, explain how to find the determinant of the matrix described in (c).

Solution:

- $P = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 4 & 0 & 1 \end{bmatrix}$

$$2. Q = \begin{bmatrix} -4 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

3. QPA

$$4. \det(QPA) = \det(Q) \det(P) \det(A) = (-4)(1)(-5) = 20.$$

□

G2. Show how to compute the determinant of the matrix

$$A = \begin{bmatrix} 1 & 3 & 0 & -1 \\ 1 & 1 & 2 & 4 \\ 1 & 1 & 1 & 3 \\ -3 & 1 & 2 & -5 \end{bmatrix}$$

Solution: Here is one possible solution, first applying a single row operation, and then performing Laplace/cofactor expansions to reduce the determinant to a linear combination of 2×2 determinants:

$$\begin{aligned} \det \begin{bmatrix} 1 & 3 & 0 & -1 \\ 1 & 1 & 2 & 4 \\ 1 & 1 & 1 & 3 \\ -3 & 1 & 2 & -5 \end{bmatrix} &= \det \begin{bmatrix} 1 & 3 & 0 & -1 \\ 0 & 0 & 1 & 1 \\ 1 & 1 & 1 & 3 \\ -3 & 1 & 2 & -5 \end{bmatrix} = (-1) \det \begin{bmatrix} 1 & 3 & -1 \\ 1 & 1 & 3 \\ -3 & 1 & -5 \end{bmatrix} + (1) \det \begin{bmatrix} 1 & 3 & 0 \\ 1 & 1 & 1 \\ -3 & 1 & 2 \end{bmatrix} \\ &= (-1) \left((1) \det \begin{bmatrix} 1 & 3 \\ 1 & -5 \end{bmatrix} - (1) \det \begin{bmatrix} 3 & -1 \\ 1 & -5 \end{bmatrix} + (-3) \det \begin{bmatrix} 3 & -1 \\ 1 & 3 \end{bmatrix} \right) + \\ &\quad (1) \left((1) \det \begin{bmatrix} 1 & 1 \\ 1 & 2 \end{bmatrix} - (3) \det \begin{bmatrix} 1 & 1 \\ -3 & 2 \end{bmatrix} \right) \\ &= (-1)(-8 + 14 - 30) + (1)(1 - 15) \\ &= 10 \end{aligned}$$

Here is another possible solution, using row and column operations to first reduce the determinant to a 3×3 matrix and then applying a formula:

$$\begin{aligned} \det \begin{bmatrix} 1 & 3 & 0 & -1 \\ 1 & 1 & 2 & 4 \\ 1 & 1 & 1 & 3 \\ -3 & 1 & 2 & -5 \end{bmatrix} &= \det \begin{bmatrix} 1 & 3 & 0 & -1 \\ 0 & 0 & 1 & 1 \\ 1 & 1 & 1 & 3 \\ -3 & 1 & 2 & -5 \end{bmatrix} = \det \begin{bmatrix} 1 & 3 & 0 & -1 \\ 0 & 0 & 1 & 0 \\ 1 & 1 & 1 & 2 \\ -3 & 1 & 2 & -7 \end{bmatrix} \\ &= -\det \begin{bmatrix} 1 & 3 & 0 & -1 \\ 1 & 1 & 1 & 2 \\ 0 & 0 & 1 & 0 \\ -3 & 1 & 2 & -7 \end{bmatrix} = -\det \begin{bmatrix} 1 & 3 & -1 \\ 1 & 1 & 2 \\ -3 & 1 & -7 \end{bmatrix} \\ &= -((-7 - 18 - 1) - (3 + 2 - 21)) \\ &= 10 \end{aligned}$$

□

G3. Explain how to find the eigenvalues of the matrix $\begin{bmatrix} -2 & -2 \\ 10 & 7 \end{bmatrix}$.

Solution: Compute the characteristic polynomial:

$$\det(A - \lambda I) = \det \begin{bmatrix} -2 - \lambda & -2 \\ 10 & 7 - \lambda \end{bmatrix} = (-2 - \lambda)(7 - \lambda) + 20 = \lambda^2 - 5\lambda + 6 = (\lambda - 2)(\lambda - 3)$$

The eigenvalues are the roots of the characteristic polynomial, namely 2 and 3.

□

G4. Explain how to find a basis for the eigenspace associated to the eigenvalue 3 in the matrix

$$\begin{bmatrix} -7 & -8 & 2 \\ 8 & 9 & -1 \\ \frac{13}{2} & 5 & 2 \end{bmatrix}.$$

Solution: The eigenspace associated to 3 is the kernel of $A - 3I$, so we compute

$$\text{RREF}(A - 3I) = \text{RREF} \begin{bmatrix} -7 - 3 & -8 & 2 \\ 8 & 9 - 3 & -1 \\ \frac{13}{2} & 5 & 2 - 3 \end{bmatrix} = \text{RREF} \begin{bmatrix} -10 & -8 & 2 \\ 8 & 6 & -1 \\ \frac{13}{2} & 5 & -1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & -\frac{3}{2} \\ 0 & 0 & 0 \end{bmatrix}.$$

Thus we see the kernel is

$$\left\{ \begin{bmatrix} -a \\ \frac{3}{2}a \\ a \end{bmatrix} \mid a \in \mathbb{R} \right\}$$

which has a basis of $\left\{ \begin{bmatrix} -1 \\ \frac{3}{2} \\ 1 \end{bmatrix} \right\}$.

□